

TECHNICAL WORKING PAPER SERIES

SOLVING LARGE SCALE RATIONAL
EXPECTATIONS MODELS

Jess Gaspar
Kenneth L. Judd

Technical Working Paper 207

NATIONAL BUREAU OF ECONOMIC RESEARCH
1050 Massachusetts Avenue
Cambridge, MA 02138
February 1997

We acknowledge the support of the National Science Foundation. This paper is part of NBER's research program in Public Economics. Any opinions expressed are those of the authors and not those of the National Bureau of Economic Research.

© 1997 by Jess Gaspar and Kenneth L. Judd. All rights reserved. Short sections of text, not to exceed two paragraphs, may be quoted without explicit permission provided that full credit, including © notice, is given to the source.

Solving Large Scale Rational Expectations Models
Jess Gaspar and Kenneth L. Judd
NBER Technical Working Paper No. 207
February 1997
JEL No. C68
Public Economics

ABSTRACT

We explore alternative approaches to numerical solutions of large rational expectations models. We discuss and compare several current alternatives, focussing on the tradeoffs in accuracy, space, and speed. The models range from representative agent models with many goods and capital stocks, to models of heterogeneous agents with complete or incomplete asset markets. The methods discussed include perturbation and projection methods. We show that these methods are capable of analyzing moderately large models even when we use only elementary, general purpose numerical methods.

Jess Gaspar
Department of Economics
Stanford University
Stanford, CA 94305
jgaspar@leland.stanford.edu

Kenneth L. Judd
Hoover Institution
Stanford University
Stanford, CA 94305
and NBER
judd@hoover.stanford.edu

The study of macroeconomic dynamics has become substantially more sophisticated over the past twenty years. The first models were simple models with one commodity, serving as both the consumption and investment good, and with one type of agent as in the “representative agent” models, or very similar agents, as in two-period overlapping generations models where agents differed only as to age. While these simple models provided insights on many issues, their limitations are becoming more apparent and we now want to move beyond them. In particular, it is desirable to add multiple goods, multiple capital stocks, heterogeneous agents, multiple assets, heterogeneous taxation, externalities, imperfect competition, and asymmetric information to the conventional models. Outside of a few special analyses, there are no examples of general dynamic, multi-agent, multi-good, stochastic, rational expectations models which handle high- and low-frequency economic movements, both of which are critical for understanding data and analyzing policies. We need to turn to numerical methods to solve these models. The methods discussed in this paper move us a step in that direction.

In this paper we address the problem of computing equilibria of large rational expectations models. We show how extensions of methods used in the simple models can be used to solve more complex models. The fact that the general models considered below can be solved numerically is obvious since one can take any of the many methods which have been developed and apply them to these models. That is not an interesting observation since most of these methods’ time and space requirements, such as discrete state space dynamic programming, would make them impractical to solve on any computer. The problem is to develop methods which can solve larger models in reasonable time using accessible resources.

This paper focusses on the application of both perturbation and projection methods to multidimensional dynamic models. In previous papers, Judd and Guu(1993, forthcoming) examined perturbation methods which go beyond the standard linearization method, and Judd(1992) examined projection methods for solving rational expectations models. Both papers focussed on applications to a representative agent, single good model. The reader is referred to these papers and their mathematical sources for key definitions and introductions to these methods. In this paper, we will outline how these methods can be adapted to handle multi-agent generalizations; in

the case of projection methods we stay with the single-good, single-asset assumptions of the representative agent model, but the perturbation analysis includes multiple goods and multiple capital stocks. A theme of Judd(1992) is that there are many ways to solve rational expectations models, the best way depending on a variety of considerations. In this paper we continue that theme attempting to determine the relationship between the most efficient method and the characteristics of the problem. We give examples of various methods, their time and space needs, and the accuracy of the results.

Our results point to a “production possibilities frontier” of methods for solving rational expectations models. If the model is stable with an ergodic distribution concentrated in the neighborhood of some deterministic steady state, then the Taylor series expansion produced by perturbation methods will likely produce a good approximation and, according to several existing examples, produce solutions which are reliable for a nontrivial region. If the nonlinearities are substantial and/or we need a more global approximation, we will likely use some kind of projection method¹.

The results below give some guidance as to which methods are most efficient in terms of the trade-off between accuracy and running time. They indicate that perturbation methods are the most efficient methods when they work, and the only feasible method for the largest problems. Among the projection methods, the best choice for moderate-size models appears to use complete polynomial approximations of the unknown functions, and a version of successive approximation introduced by Miranda and Helmberger(1988) to solve for the unknown coefficients. The dominance of complete polynomial representation is not surprising, but the success of successive approximation is a bit surprising since it is not even locally convergent in general. However, its considerable computational advantages over Newton methods for solving nonlinear equations combined with our favorable experience (below and in earlier papers) indicates that it should generally be considered. One should keep in mind, however, that all the models we examine below have strong mean reversion properties in equilibrium, a property which may help explain the success of successive approximation. Because these experiments indicate that successive approximation is often valuable, future work should focus on exactly when it is stable.

While these conclusions are based on the analysis of simple examples, we feel that

this experience is likely to be robust to other models and that its conclusions can serve as a reasonable guide to choosing methods for solving large models. Hopefully the message here is encouraging enough that analysts will feel free to analyze large models of their choosing.

1. NUMERICAL RATIONAL EXPECTATIONS PROBLEMS

Before we discuss various specific rational expectations models, we should point out the features which are inherent in numerical rational expectations modelling. Awareness of these features will help us form reasonable modelling goals and point in the direction of appropriate numerical methods.

In theory, rational expectations models resemble dynamic interpretations of Arrow-Debreu general equilibrium analysis. The central focus of Arrow-Debreu models are the time- and state-contingent equilibrium prices and consumption patterns. The determination of equilibrium prices is the focus of standard CGE methods. However, this is an unreasonable numerical approach in the case of rational expectations models. In the Arrow-Debreu approach to infinite-horizon dynamic stochastic models, the price of each good is contingent on the date at which it appears and the entire history of the exogenous shocks up to that date. In the case of discrete-time and infinite-horizon, if there are exogenous shocks in each period and the support of these shocks contains at least two points, the number of distinct histories in an infinite-horizon model has the cardinality of the continuum, implying that the number of distinct prices is also the size of the continuum and that the conventional CGE approach is infeasible.

These problems can be avoided if the structural features of the economic model are stationary, that is, do not depend on calendar time, and if we focus on stationary rational expectations equilibria of those stationary infinite horizon models. Such equilibria can be expressed in terms of decision rules for the agents (see Stokey and Lucas(1989) for a formal discussion of recursive modelling), and the numerical approach is to compute approximations of these unknown functions. For example, a simple discrete-time problem explored in the computational literature is

$$\begin{aligned} \max_c \quad & E \{ \sum_{t=0}^{\infty} \beta^t u(c_t) \} \\ & k_{t+1} = F(k_t, \theta_t) - c_t \\ & \ln \theta_{t+1} = \rho \ln \theta_t + \epsilon_{t+1} \end{aligned} \tag{1}$$

where k_t is the beginning-of-period capital stock, θ_t is a stationary AR(1) multiplicative productivity parameter (the productivity shocks $\epsilon_t \sim N(0, \sigma^2)$ are independent), and $F(k, \theta)$ is the gross (that is, net production plus initial capital stock) production function. In this problem, both k and θ are needed for a sufficient description of the state. Hence, consumption is a function of both k and θ , $C(k, \theta)$, and the Euler equation is

$$u'(C(k, \theta)) = \beta E \{u'(C(F(k, \theta) - C(k, \theta), \tilde{\theta}))F_k(F(k, \theta) - C(k, \theta), \tilde{\theta}) | \theta\} \quad (2)$$

This is the problem investigated in the Taylor–Uhlig symposium (1990) and in Judd(1992). While we will focus on generalizations of (2), our results apply also to the commodity problems studied in earlier computational rational expectations analyses by Gustafson(1958), Wright and Williams(1984, 1991), and Miranda and Helmberger(1988).

At first, focussing on the functional equation in (2) appears to be no better since we move from the infinite-dimensional space of contingent prices to the infinite-dimensional space of decision rules. There is improvement, however, since there are good finite-dimensional approximations of the equilibrium policy functions such as $C(k, \theta)$ when they are “reasonable” functions, whereas we know of no such way to approximate the continuum of contingent prices. Numerical rational expectations methods, beginning with Gustafson(1958), therefore focus on finite-dimensional approximations of policy functions and other important functions. The finite-dimensional approximations typically parameterize the unknown policy function² and restrict it to lie in some finite-dimensional space, as in

$$\hat{C}(k, \theta) = \sum_{i=0}^n a_i \phi_i(k, \theta)$$

where the ϕ_i comprise a basis for all candidate functions. We must first make a choice of basis, and then solve for the unknown coefficients, a . There are several ways to fix these unknown coefficients. Projection methods fix a by solving a set of projection equations which try to identify a which will nearly solve (2). Perturbation methods construct a Taylor series expansion around a point (k_0, θ_0) , such as

$$\hat{C}(k, \theta) = \sum_{i=0}^n \sum_{j=0}^n a_{ij} (k - k_0)^i (\theta - \theta_0)^j$$

and use implicit function theorems to identify the undertermined a_{ij} coefficients.

The various approaches to solving rational expectations models differ in three basic ways; first, in the choice of finite-dimensional approximations to functions, second, in the way the expectation in (2) is computed, and, third, in the method used to find an approximate solution. The work discussed below touches on two of the three critical elements — the method used to approximate C and the method for solving the identifying conditions. In this paper we will focus on various combinations of approximation and solution methods which appear to be promising in the context of large rational expectations models.

We will discuss both perturbation and projection methods. They share much in common; see Judd(1996) for an extended discussion of this observation. We will focus on applications of perturbation methods to continuous-time models and applications of projection methods to discrete-time models. The dynamic economics literature bounces between continuous- and discrete-time applications depending on whether one builds on the Brock and Turnovsky(1981) analysis of dynamic macroeconomic equilibrium, or the Brock and Mirman(1972) stochastic growth model. There is no substantive economic difference since the discrete time unit can be made arbitrarily small. We make our choices to focus on the simple cases; both methods are applicable to both types of models, but perturbation analysis of discrete-time models is much more complex notationally, and projection analysis of continuous-time models would require us to introduce functional analytic material of little interest to the intended reader. We leave these developments to future work.

2. PERTURBATION METHODS

We will first explore an example of perturbation methods applied to the canonical continuous-time, stable optimal control problem. We will not define perturbation in general here, nor will we present the mathematical foundations which justify the formal, algebraic manipulations presented below; see Judd(forthcoming, 1996) and Bensoussan(1988) for general discussions and for citations of the formal development of perturbation methods for control problems. Here we will take a “linearize around the steady-state” approach to large rational expectations models and extend it, via perturbation methods, to include nonlinear terms of the Taylor series expansion. While the linearization approach is common, general perturbation methods have been used

only to a limited extent in economics. There are several reasons for this. First, there is little agreement in the literature³ as to what constitutes “linearization.” Second, even those who “linearize” deterministic models correctly⁴ generally fail to compute the true first-order Taylor series approximation when they approximate stochastic models; we make this precise below. Third, there is very limited mention in the economics literature of the higher-order terms in the expansion implicitly being constructed. In fact, Marcet(1995) states that “perturbation methods of order higher than one are considerably more complicated⁵ than the traditional linear-quadratic case ...” Below we show that while computing the traditional linear-quadratic approximation involves solving a Riccati-like equation, computing those dreaded higher-order terms involves solving only linear equations, and, therefore, that computing some higher-order terms is in fact less demanding computationally than computing the initial linear term.

High-order expansions are useful for two reasons. First, higher-order terms are necessary for analyzing the first-order properties of many aspects of the model; an example of this would be the cyclical properties of risk premia which clearly involves how the risk aversion coefficient moves over time, a third-order property of utility. The second reason, and the focus of this section, is that Taylor series expansions which include nonlinear terms may provide good approximations over a larger region than the linear approximation would. The fact that much effort has been extended to compute nonlinear solutions to dynamic growth models, such as in Taylor and Uhlig(1990), indicates that we do not believe that linear approximations are always adequate.

In this section, we present the standard mathematical procedure for linearizing a model around a stable steady state⁶, for computing higher-order terms in the expansion, and for computing the extra terms necessary for an asymptotically valid expansion in stochastic models. We will break down the method into the relevant computational steps, and evaluate the computational cost of large models. In this paper we focus on the mechanics of computing Taylor series expansions for large, general dynamic programming problems and the associated computational demands. Future work will examine the quality of the approximation in various specific contexts, and on generalizing the analysis to the case of competitive equilibrium.

Before continuing, we should warn the reader of the nontrivial notational chal-

lenge which awaits him in the sections below. After being introduced to tensor notation and its application to multivariate stochastic control, the reader may decide that this approach is far too burdensome to be of value. If one had to go through these manipulations for each and every application, we might agree. Fortunately, all of the algebra discussed below can be automated. Furthermore, the authors are writing user-friendly programs which will take specifications of tastes and technology (represented by some user-written subroutines) and will automatically perform *all* the necessary computations, including derivatives, and produce the Taylor series approximation discussed below. This will relieve the user of executing all the algebra we discuss below. This software will be available at the anonymous ftp site at bucky.stanford.edu. The presentation below is meant to familiarize the reader with the mathematical structure of the problem. To help interested readers better understand the algebraic details, we have also worked out a two-dimensional example in complete detail. It would consume about twenty pages in this paper, so we also will put that at bucky.stanford.edu.

2.1. Uses of Taylor Series Approximations. To motivate the following computations, we next indicate what we can do with these Taylor series approximations. First, the focus of this paper, we can use them to serve as global (or, more precisely, nonlocal) approximations. This may seem inappropriate since the Taylor series construction is only local⁷. However, the global accuracy experiments conducted so far (for example, see Judd and Guu (1993, forthcoming) indicate that these local methods do well in nontrivial neighborhoods of the steady state in dynamic economic problems as long as the utility and production functions are analytic in a neighborhood of the deterministic steady state. Second, we can produce linear theories of any interesting economic phenomenon⁸. Linear approximations of policy functions will produce linear theories about equilibrium choices, such as consumption, investment, and output. However, if we want to produce a linear theory for other aspects of equilibrium we will often need higher order terms. A linear theory for the movement of the equity risk premium over the business cycle, or a similar linear theory of the term structure of interest rates, requires higher-order Taylor expansions of the equilibrium decision rules.

Third, these approximations are the first step in potential empirical procedures since a linear theory of an economic quantity can be used to compare theory and data. For example, Magill(1977) showed how to use linear approximations of a stochastic growth model to compute the implied spectrum for consumption, and suggested that comparing such theoretical spectra with empirical spectra can be a useful empirical approach to business cycle investigations. Later work by Kydland and Prescott(1982) used Magill's approach to compare data and a particular stochastic growth model. The Taylor series we compute below can be similarly used to compare theory and data since these expansions can also be used to compute locally valid approximations of likelihood functions and their derivatives. Fourth, these perturbation methods have been used in policy evaluation exercises. See Judd(1996) for a more extensive discussion of these topics.

2.2. The General Dynamic Optimization Problem. For the purposes of this discussion, it is advantageous to discuss the continuous-time dynamic programming case⁹. We could use the equivalence between competitive equilibrium and Pareto efficiency to derive approximations to competitive equilibrium decision rules. These methods can handle distortions, but with some added complexity which the authors leave for future work. In general, if we have many agents, many goods, and many capital stocks, including "stocks" in the utility functions, the dynamic optimization problem is

$$\begin{aligned} V(x_0) \equiv \max_{u(t)} \int_0^{\infty} e^{-\rho t} \pi(x, u) dt \\ \dot{x} = f(x, u) \\ x(0) = x_0 \end{aligned} \quad (3)$$

where $x \in R^n$ is the state vector, $u \in R^m$ is the vector of controls, $f(x, u)$ is the law of motion, and $\pi(x, u)$ is a concave social welfare function. This problem has the Hamiltonian $H(x, \lambda, u) = \pi(x, u) + \lambda f(x, u)$, implying the differential-algebraic system.

$$\begin{aligned} \dot{x} &= f(x, u) & &= H_\lambda \\ \dot{\lambda} &= \rho\lambda - (\lambda f_x(x, u) + \pi_x) & &= \rho\lambda - H_x \\ 0 &= \pi_u(x, u) + \lambda f_u(x, u) & &= H_u \end{aligned} \quad (4)$$

When we substitute the control law $\mathcal{U}(x, \lambda)$, implicitly defined by $0 = \pi_u(x, \mathcal{U}(x, \lambda)) + \lambda f_u(x, \mathcal{U}(x, \lambda))$, into (4) we arrive at the dynamic system

$$\begin{aligned}\dot{x} &= f(x, \mathcal{U}(x, \lambda)) \\ \dot{\lambda} &= \rho\lambda - (\lambda f_x(x, \mathcal{U}(x, \lambda)) + \pi_x(x, \mathcal{U}(x, \lambda)))\end{aligned}\quad (5)$$

Furthermore, we are only interested in asymptotically bounded solutions to this equation with the initial condition $x(0) = x_0$. This is the typical kind of deterministic dynamical system we begin with in our linearization exercises.

2.3. Local Dynamics. To linearize systems such as (5), one invokes basic ordinary differential equation theory. Let $Z = \begin{pmatrix} x \\ \lambda \end{pmatrix}$. Suppose we have a dynamic system

$$\dot{Z} = g(Z) \quad (6)$$

with a stationary point at Z^* ; that is, $g(Z^*) = 0$. Then the local behavior of (6) for Z near Z^* is linearly approximated by the linear system

$$\dot{z} = A z \quad (7)$$

where $A = g_Z(Z^*)$ and $z \equiv Z - Z^*$. The solution to (7) is $z(t) = e^{At} z_0$.

In the terminology of linear rational expectations models, as in Blanchard and Kahn[3], the vector x contains the predetermined variables and λ contains the variables with free values at $t = 0$. Suppose that there is a stationary point at $Z^* = \begin{pmatrix} x^* \\ \lambda^* \end{pmatrix}$. Then the local behavior of the system is linearly approximated by (7) and the solution to the linear approximation is $z(t) = e^{At} \begin{pmatrix} x(0) - x^* \\ \lambda(0) - \lambda^* \end{pmatrix}$, where $x(0) = x_0$ is a given initial condition and $\lambda(0)$ is chosen to keep $z(t)$ bounded asymptotically. Let $\Lambda(x_0)$ be the set of all possible values for the free variables in λ which together with the predetermined variables being equal to x_0 will imply a bounded path for $z(t)$. $\Lambda(x_0)$ may be a single value or a set of values. We will assume that it is single-valued, the case of determinacy. We will see where that is necessary in our calculations.

To compute $\Lambda(x_0)$, we just apply standard linear algebra. We form the Jordan decomposition of $A = N^{-1}DN$ where $D = \begin{pmatrix} D_1 & 0 \\ 0 & D_2 \end{pmatrix}$ with D_1 having all the stable eigenvalues of A (that is, the eigenvalues with negative real parts) and D_2 having the

unstable eigenvalues, and $N = \begin{pmatrix} N_{11} & N_{12} \\ N_{21} & N_{22} \end{pmatrix}$ breaks N into equal size blocks. If the number of stable eigenvalues equals the length of x_0 , stability of the solution to (7) implies

$$\Lambda(x_0) - \lambda^* = -N_{22}^{-1}N_{21}(x_0 - x^*) \quad (8)$$

We can apply this same approach to discrete-time systems. Suppose that we have a system, $Z_{t+1} = g(Z_t)$, with a steady state defined by $Z^* = g(Z^*)$. Define $A = g_Z(Z^*)$ and $z \equiv Z - Z^*$. Then the equation for z becomes $z_{t+1} = Az_t$, the solution of which is $z_t = A^t z_0$, which in turn can be analyzed in the same Jordan decomposition fashion with the distinction that now the stable eigenvalues are those with modulus less than one. Unfortunately, the general discrete-time case is more complex than $Z_{t+1} = g(Z_t)$; see Judd and Gaspar(1997) for a complete treatment of the discrete-time case.

2.4. Higher-Order Approximations. We will next discuss the computational demands of computing higher-order terms to the multivariate Taylor series approximation based at the deterministic steady state. This may seem formidable. To solve the first-order terms we had to solve an eigenvalue-eigenvector problem. In fact, if one were to look at the details, one would find that the problem is very similar to a Riccati equation, that is, a quadratic matrix equation. When discussing this problem with macroeconomists, we have often heard the conjecture that computing higher-order terms would require solving higher order matrix polynomial problems. The basic fact shown below is that all the higher-order terms of the Taylor series expansion, even in the stochastic multidimensional case, are solutions to linear problems once one computes the first-order terms. This implies that the higher-order terms are *easy* to compute. Initial experiments indicate that they are also good approximations well beyond the steady state values. These procedures have not been exploited, but can be obviously applied to problems in the real business cycle, finance, public finance, and dynamic general equilibrium literatures.

Tensor Notation. To deal with the notational problems which arise with multidimensional expressions, we extend the Einstein tensor notation, adapting it for our control theoretic problems. In general, a tensor is any indexed collection of numbers.

The second property is that we use the position and repetition of indices to indicate summation. Suppose that a_i is a collection of numbers indexed by $i = 1, \dots, n$, and that x^i is also a singly indexed collection of real numbers. Then

$$a_i x^i \equiv \sum_i a_i x^i$$

In this way we eliminate the Σ symbol in expressing vector products. Similarly, suppose that a_{ij} is a collection of numbers indexed by $i, j = 1, n$, and that x^i and y^j are singly indexed collections of real numbers. Then

$$a_{ij} x^i y^j \equiv \sum_i \sum_j a_{ij} x^i y^j$$

In this way we again eliminate the Σ symbols. The general rule shall be that we eliminate Σ symbols by understanding that, in a product, if an index appears as both a subscript and a superscript, then we sum over it. If we think of a_{ij} as a matrix, x_i as a row vector, and y^j as a column vector, then the product $x_i y^i$ represents the inner product of the vectors x and y , and $a_{ij} x_i y^j$ is the quadratic form of the matrix a with the vectors x and y . We can also form new indexed collections of numbers from products. For example, the product $a_{ij} x_i$ can be thought of as a singly indexed collection of numbers, z_j . Using our vector analogy above, z_j can also be thought of as a row vector. While the analogies with matrices and vectors are useful, one should not focus on them because we will be constructing more complex collections of real numbers which are neither vectors nor matrices.

As long as indices are not the same, arbitrary products are allowed. For example, $x_i y_j$ is the doubly indexed set of numbers, b_{ij} , where the (i, j) term equals the product of x_i and y_j ; b_{ij} is the *outer product* of x_i and y_j . Also, the "sum over repeated indices" rule even applies within a single term. For example, $a_i^i \equiv \sum_i a_i^i$; if we think of a as a matrix, then a_i^i is the trace of a . Similarly, a_{ii} is also the trace of the tensor a_{ij} .

We will also vary the notation in order to distinguish between an argument of a vector and a derivative, and between states and controls in an efficient fashion. In the notation below, superscripts will refer to different components of a vector-valued function, whereas subscripts will refer to derivatives of those component functions. Furthermore, in order to distinguish between states and controls, we shall let Roman letters, i, j, k, ℓ, \dots , index states, and Greek letters, $\alpha, \beta, \gamma, \dots$, index controls. There-

fore, if $f(x, u)$ is a vector-valued function of the state variables x and the controls u , then f is the column vector

$$f = \begin{pmatrix} f^1(x, u) \\ f^2(x, u) \\ \vdots \\ f^n(x, u) \end{pmatrix}$$

whose i 'th component function is denoted f^i . Its derivatives with respect to the state variables are represented by the tensor

$$f_i^j(x, u) \equiv \frac{\partial f^j}{\partial x_i}(x, u)$$

and its derivatives with respect to the control variables are represented by the tensor

$$f_\alpha^j(x, u) \equiv \frac{\partial f^j}{\partial u_\alpha}(x, u)$$

We will frequently drop the argument (x, u) as long as it can be understood from context.

So far, all of this looks familiar since it is just a way of rewriting standard matrix and vector operations. However, triply indexed collections of numbers, such as $a_{ij\ell}$, are also tensors, and arise naturally in multivariate calculus. For example, Taylor's theorem for $g : R^n \rightarrow R$ at $z = 0$ is normally written as

$$\begin{aligned} f(z) \sim & f(0) + \sum_{i=1}^n \frac{\partial f}{\partial z_i}(0)z_i + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2 f}{\partial z_i \partial z_j}(0)z_i z_j \\ & + \frac{1}{6} \sum_{i=1}^n \sum_{j=1}^n \sum_{\ell=1}^n \frac{\partial^3 f}{\partial z_i \partial z_j \partial z_\ell}(0)z_i z_j z_\ell + \dots \end{aligned}$$

whereas by using tensor notation it can be written as

$$f(z) \sim f(0) + f_i(0)z^i + \frac{1}{2}f_{ij}(0)z^i z^j + \frac{1}{6}f_{ij\ell}(0)z^i z^j z^\ell + \dots$$

If we drop the arguments of f , and understand that f and its derivatives are evaluated at $x = 0$, the Taylor expansion can be written as

$$f(z) \sim f + f_i z^i + \frac{1}{2}f_{ij} z^i z^j + \frac{1}{6}f_{ij\ell} z^i z^j z^\ell + \dots$$

This more compact expression is a considerable improvement over the conventional notation with the extraneous summation symbols and the clumsy partial derivative notation. With this notation, we will be able to more clearly see the structure of our problem.

Computing Taylor Series Expansions for Optimal Control Problems.

The general deterministic problem (3) includes an arbitrary complete market dynamic equilibrium with multiple consumption goods (represented by components of u), multiple capital stocks (represented by components of x) and multiple agents whose aggregate utility function is represented by $\pi(x, u)$. We will now proceed with a dynamic programming approach to this analysis. The Bellman equation for the value function, $V(x)$, is

$$0 = \max_u \pi(x, u) + V_i(x) f^i(x, u) - \rho V(x) \quad (9)$$

where, recall, π is the payoff flow and f is the law of motion for the state x . The first-order condition with respect to u^α , $\alpha = 1, \dots, m$, is

$$0 = \pi_\alpha(x, u) + V_i(x) f_\alpha^i(x, u) \quad (10)$$

Equations (10) implicitly define the optimal control, $u = U(x)$, and imply the system

$$0 = \pi_\alpha(x, U(x)) + V_i(x) f_\alpha^i(x, U(x)) \quad (11)$$

In combination, we have the system

$$0 = \pi(x, U(x)) + V_i(x) f^i(x, U(x)) - \rho V(x) \quad (12)$$

$$0 = \pi_\alpha(x, U(x)) + V_i(x) f_\alpha^i(x, U(x)) \quad (13)$$

which defines the value function, $V(x)$, and the policy function, $U(x)$.

Our objective here is to solve for both the value function and the policy function. In fact, we are going to compute Taylor series expansions

$$\begin{aligned} V(x) &= V(x^0) + V_i(x - x^0)^i + \frac{1}{2} V_{ij}(x - x^0)^i (x - x^0)^j \\ &\quad + \frac{1}{3!} V_{ij\ell}(x - x^0)^i (x - x^0)^j (x - x^0)^\ell + \dots \\ U^\alpha(x) &= U^\alpha(x^0) + U_i^\alpha(x - x^0)^i + \frac{1}{2} U_{ij}^\alpha(x - x^0)^i (x - x^0)^j \\ &\quad + \frac{1}{3!} U_{ij\ell}^\alpha(x - x^0)^i (x - x^0)^j (x - x^0)^\ell + \dots \end{aligned} \quad (14)$$

These polynomials are asymptotically valid approximations if the error converges to zero at a higher degree than the order of the polynomial. For example, the linear approximation $V(x^0) + V_i(x - x^0)^i$ has an error which is quadratic in the components of $(x - x^0)$, which means that as x converges to x^0 the error goes to zero quadratically.

To compute these U_i^α , U_{ij}^α , U_{ijl}^α , V_i , V_{ij} , and V_{ijl} coefficients we just differentiate the underlying system, (12,13), with respect to the x_i , and solve for the undetermined coefficients. If we differentiate (12) with respect to x_j and use the envelope theorem we find

$$\rho V_j = \pi_j + V_{ij} f^i + V_i f_j^i \quad (15)$$

In order to keep down the clutter, we often drop the arguments of π , V , and U , and their derivatives when they are the same as in the basic system, (12,13) and are clear from context.

The steady state values for x , u , and V_i are determined by the conditions

$$\begin{aligned} 0 &= f^i(u, x) \\ 0 &= \pi_\alpha(u, x) + V_i(x) f_\alpha^i(u, x) \\ \rho V_j(x) &= \pi_j(u, x) + V_{ij}(x) f^i(u, x) + V_i(x) f_j^i(u, x) \end{aligned} \quad (16)$$

which yield the steady-state quantities u^{ss} , x^{ss} , $V_j(x^{ss})$. Note that the $V_j(x^{ss})$ values are the linear coefficients in the expansion of V in (14), and knowing the steady state will also yield $V(x^{ss})$ and $U(x^{ss})$, two more terms in (14).

We next compute the V_{ij} and U_j^β terms, and then compute many high-order derivatives. We assume that these derivatives exist, which leads us to assume that all production and utility functions are C^∞ . We *assume* the differentiability of the value and policy functions in the neighborhood of the deterministic steady state. For deterministic problems, this is usually proven by applying theorems about the smooth dependence of differential equation solutions on parameters, and Fleming(1971) deals with the stochastic problem. We will proceed under the assumption that the indicated derivatives exist.

If we differentiate (13) with respect to the x_j we find

$$0 = \pi_{\alpha j} + \pi_{\alpha \gamma} U_j^\gamma + V_{ij} f_\alpha^i + V_i (f_{\alpha j}^i + f_{\alpha \gamma}^i U_j^\gamma) \quad (17)$$

Note that (17) is a system of conditions, one for each αj pair. In this case, we can

express U_j^γ in terms of the derivatives of V

$$U_j^\gamma = -(\pi_{\alpha\gamma} + V_i f_{\alpha\gamma}^i)^{-1} (\pi_{\alpha j} + V_{ij} f_\alpha^i + V_i f_{\alpha j}^i) \quad (18)$$

where $(\pi_{\alpha\gamma} + V_i f_{\alpha\gamma}^i)^{-1}$ denotes the inverse tensor (matrix). Differentiating (15) with respect to x_ℓ implies

$$\begin{aligned} \rho V_{j\ell} &= \pi_{j\ell} + \pi_{j\gamma} U_\ell^\gamma + V_{ij\ell} f^i + V_{ij} (f_\ell^i + f_\gamma^i U_\ell^\gamma) \\ &\quad + V_{i\ell} f_j^i + V_i (f_{j\ell}^i + f_{j\gamma}^i U_\ell^\gamma) \end{aligned} \quad (19)$$

Substituting (18) into (19) yields

$$\begin{aligned} \rho V_{j\ell} &= \pi_{j\ell} + V_{ij\ell} f^i + V_{ij} f_\ell^i + V_{i\ell} f_j^i + V_i f_{j\ell}^i \\ &\quad - (\pi_{j\gamma} + V_{ij} f_\gamma^i + V_i f_{j\gamma}^i) (\pi_{\alpha\gamma} + V_i f_{\alpha\gamma}^i)^{-1} (\pi_{\alpha\ell} + V_{i\ell} f_\alpha^i + V_i f_{\alpha\ell}^i) \end{aligned} \quad (20)$$

The system of equations in (20) hold at each state x . If we evaluate (20) at the steady state, then $f^i = 0$ and (20) becomes the Riccati-like equation

$$\begin{aligned} \rho V_{j\ell} &= \pi_{j\ell} + V_{ij} f_\ell^i + V_{i\ell} f_j^i + V_i f_{j\ell}^i \\ &\quad - (\pi_{j\gamma} + V_{ij} f_\gamma^i + V_i f_{j\gamma}^i) (\pi_{\alpha\gamma} + V_i f_{\alpha\gamma}^i)^{-1} (\pi_{\alpha\ell} + V_{i\ell} f_\alpha^i + V_i f_{\alpha\ell}^i) \end{aligned} \quad (21)$$

Solving the Riccati equation at the steady state yields the steady state values of $V_{j\ell}$ and, through (18), the steady-state values of U_ℓ^γ . But we already have the solutions for the steady-state values of U_ℓ^γ because the U_ℓ^γ tensor can be derived from the definition of $\mathcal{U}(x, \lambda)$ and the linear approximation computed in (8). This approximation was computed directly from eigenvalue decomposition methods. Therefore, we use the earlier approach to compute the elements of the tensor (matrix) U_ℓ^γ . While we have not accomplished anything new at this point, this lays the foundation for the higher-order terms.

We now find out how easy it is to compute the higher-order terms. Differentiating (19) with respect to x_m and imposing the steady state condition $f^i = 0$ implies the following equation for the steady state values of $V_{i\ell m}$ and $U_{\ell m}^\beta$:

$$\begin{aligned} \rho V_{j\ell m} &= \pi_{j\ell m} + \pi_{j\ell\gamma} U_m^\gamma + \pi_{j\gamma m} U_\ell^\gamma + \pi_{j\gamma\delta} U_m^\delta U_\ell^\gamma + \pi_{j\gamma} U_{\ell m}^\gamma \\ &\quad + V_{ij\ell} (f_m^i + f_\gamma^i U_m^\gamma) + V_{ijm} (f_\ell^i + f_\gamma^i U_\ell^\gamma) \\ &\quad + V_{ij} (f_{\ell m}^i + f_{\ell\gamma}^i U_m^\gamma + f_{\gamma m}^i U_\ell^\gamma + f_{\gamma\delta}^i U_\ell^\delta U_m^\delta + f_\gamma^i U_{\ell m}^\gamma) \\ &\quad + V_{i\ell m} f_j^i + V_{i\ell} (f_{j\gamma}^i U_m^\gamma + f_{jm}^i) + V_{im} (f_{j\ell}^i + f_{j\gamma}^i U_\ell^\gamma) \\ &\quad + V_i (f_{j\ell m}^i + f_{j\ell\gamma}^i U_m^\gamma + f_{j\gamma m}^i U_\ell^\gamma + f_{j\gamma\delta}^i U_\ell^\delta U_m^\delta + f_{j\gamma}^i U_{\ell m}^\gamma) \end{aligned} \quad (22)$$

When we rewrite (17) as

$$0 = (\pi_{\alpha\beta} + V_i f_{\alpha\beta}^i) U_\ell^\beta + (\pi_{\alpha\ell} + V_{i\ell} f_\alpha^i + V_i f_{\alpha\ell}^i)$$

and differentiate this expression with respect to x_m , and impose the steady state condition we find

$$\begin{aligned} 0 = & (\pi_{\alpha\beta} + V_i f_{\alpha\beta}^i) U_{\ell m}^\beta \\ & + (\pi_{\alpha\beta\gamma} U_m^\gamma + \pi_{\alpha\beta m} + V_{im} f_{\alpha\beta}^i + V_i f_{\alpha\beta m}^i + V_i f_{\alpha\beta\gamma}^i U_m^\gamma) U_\ell^\beta \\ & + \pi_{\alpha\ell m} + \pi_{\alpha\ell\gamma} U_m^\gamma + V_{i\ell m} f_\alpha^i + V_{i\ell} (f_{\alpha m}^i + f_{\alpha\gamma}^i U_m^\gamma) \\ & + V_{im} f_{\alpha\ell}^i + V_i (f_{\alpha\ell m}^i + f_{\alpha\ell\gamma}^i U_m^\gamma) \end{aligned} \quad (23)$$

We have now reached an important point in the analysis. At this point, we know the steady state values of the U_ℓ^β and V_{im} tensors as well as the steady state values of the derivatives of π and f which appear. We see that the steady state values of $V_{i\ell m}$ and $U_{\ell m}^\beta$ appear *linearly* in (23,22) and that they are the only unknowns. The system is not cubic nor quadratic, but linear in these unknowns.

The second important point is that solving this linear system is easier than it initially appears. An efficient way to compute the steady state values of $V_{i\ell m}$ and $U_{\ell m}^\beta$ in (23,22) is immediately apparent. The key observation is that we can solve the system for each fixed ℓm pair, allowing us to break down the large system into a collection of smaller ones. Consider (23) for a fixed ℓm pair. The $\pi_{\alpha\beta} + V_i f_{\alpha\beta}^i$ tensor appears repeatedly for each ℓm pair. Hence, we can use the $(\pi_{\alpha\beta} + V_i f_{\alpha\beta}^i)^{-1}$ tensor (an inversion which is done just once and used for each ℓm pair) to express $U_{\ell m}^\beta$ linearly in terms of the known steady state values of various steady state derivatives and $V_{j\ell m}$. After we gather terms, it takes just two matrix multiplications for each ℓm pair to determine the coefficients of the affine representation

$$\begin{aligned} U_{\ell m}^\beta = & -(\pi_{\alpha\beta} + V_i f_{\alpha\beta}^i)^{-1} \times [(\pi_{\alpha\beta\gamma} U_m^\gamma + \pi_{\alpha\beta m} + V_{im} f_{\alpha\beta}^i + V_i f_{\alpha\beta m}^i + V_i f_{\alpha\beta\gamma}^i U_m^\gamma) U_\ell^\beta \\ & + \pi_{\alpha\ell m} + \pi_{\alpha\ell\gamma} U_m^\gamma + V_{i\ell} (f_{\alpha m}^i + f_{\alpha\gamma}^i U_m^\gamma) \\ & + V_{im} f_{\alpha\ell}^i + V_i (f_{\alpha\ell m}^i + f_{\alpha\ell\gamma}^i U_m^\gamma)] \\ & - (\pi_{\alpha\beta} + V_i f_{\alpha\beta}^i)^{-1} f_\alpha^i V_{i\ell m} \end{aligned} \quad (24)$$

For a fixed ℓm pair, these representations of the $U_{\ell m}^\beta$ can then be substituted into the steady state value of (22) to produce a system of equations linear in the $V_{j\ell m}$. Hence, we see that the second-order terms can be computed in a sequentially *linear* fashion

once the first-order terms have been computed. If there are n states and m controls, the total computational burden is the $2n \times 2n$ eigenvalue–eigenvector decomposition, one $m \times m$ inversion, $m(m - 1)/2$ evaluations of the expression in (24), and solving a linear system of size $n(n - 1)(n - 2)/6$.

The final important fact is that we can repeat this to compute the third- and higher-order terms of U in a similar sequentially linear fashion. We suspect that the reader will believe us on this point without seeing the algebraic details. We may ask how big can we go in this fashion. Clearly, the greater the dimension, the fewer higher-order terms we can add to the expansion. We will investigate feasibility issues below.

This may appear confusing and surprising since standard intuition says that higher-order terms are more difficult to compute. The basic explanation is that the initial linearization step must deal with the multiplicity which arises due to the steady state lying on both the stable and unstable manifolds, a multiplicity manifested by the multiple solutions to the Riccati-like equation which arises. This multiplicity arises since our methods uses only the first-order conditions, conditions which describe the unstable as well as the stable manifold as well as many other manifolds. Once the linear expansion term makes a choice of which manifold to follow, a task accomplished in (8), we have fixed our attention on the stable manifold and higher-order terms involve only easy linear problems.

Multisector, Stochastic Growth. We next examine the stochastic generalization of (3), which is

$$\begin{aligned} \max_{dx^i} \quad & E\left\{\int_0^\infty e^{-\rho t} \pi(x, u) dt\right\} \\ & = f^i(x, u) dt + \sqrt{2\epsilon} I dz \end{aligned} \tag{25}$$

where dz is a vector of i.i.d. white noises of unit variance, I is the variance–covariance matrix, assumed here to be a constant to reduce the notational burden, and ϵ is a parameter expressing the absolute magnitude of the variances. Again, any dynamic, complete market general equilibrium can be represented by the solution to such a problem. The solutions to this problem can be used to represent consumption allocation processes, investment and consumption processes, as well as asset price processes.

The Bellman equation to (25) is

$$0 = \max_u \pi(x, u) + V_i f^i + \epsilon V_{ii} - \rho V \quad (26)$$

and the first-order condition is again

$$0 = \pi_\alpha + V_i f_\alpha^i \quad (27)$$

Again, we are trying to compute Taylor series approximations to V and U , but here we take into account not only deviations of x from the deterministic steady state, $x^0 - x$, but also deviations from the deterministic case, as measured by ϵ . Here the expansions are

$$V(x, \epsilon) = V(x^{ss}, 0) + V_i(x^{ss}, 0)(x^i - x^{i0}) + V_\epsilon(x^{ss}, 0)\epsilon \quad (28)$$

$$+ V_{i\epsilon}(x^{ss}, 0)(x^i - x^{i0})\epsilon + \frac{1}{2}V_{ij}(x^{ss}, 0)(x^i - x^{i0})(x^j - x^{j0}) \\ + \frac{1}{2}V_{\epsilon\epsilon}(x^{ss}, 0)\epsilon^2 + \dots \quad (29)$$

$$U^\alpha(x, \epsilon) = U^\alpha(x^{ss}, 0) + U_i^\alpha(x^{ss}, 0)(x^i - x^{i0}) + U_\epsilon^\alpha(x^{ss}, 0)\epsilon \\ + U_{i\epsilon}^\alpha(x^{ss}, 0)(x^i - x^{i0})\epsilon + \frac{1}{2}U_{ij}^\alpha(x^{ss}, 0)(x^i - x^{i0})(x^j - x^{j0}) \\ + \frac{1}{2}U_{\epsilon\epsilon}^\alpha(x^{ss}, 0)\epsilon^2 + \dots \quad (30)$$

where all of the functions on the RHS are evaluated at $(x^{ss}, 0)$. The analysis of the deterministic problem produced all of the U_i^α , U_{ij}^α , $U_{ij\ell}^\alpha$, V_i , V_{ij} , and $V_{ij\ell}$ etc. coefficients. We now want to derive the $U_{i\epsilon}^\alpha$, $U_{ij\epsilon}^\alpha$, $V_{i\epsilon}$, $V_{ij\epsilon}$, $U_{i\epsilon\epsilon}^\alpha$, $U_{ij\epsilon\epsilon}^\alpha$, etc. coefficients.

Note additional terms due to the stochastic parameter ϵ . The usual approaches (such as in Kydland and Prescott, 1982, and Christiano, 1990) take a linear approximation to the deterministic, $\epsilon = 0$, model, which here is $U^\alpha(x, \epsilon) = U^\alpha(x^{ss}, 0) + U_i^\alpha(x^{ss}, 0)(x^i - x^{i0})$, and then use that linear rule as an approximate solution to a stochastic version of the model. This ignores the additional $U_\epsilon^\alpha(x^{ss}, 0)\epsilon$ term which is suggested by this Taylor series approach. Since the objective of all these methods is to approximate the value of u at state $x \neq x^{ss}$ in a model where $\epsilon \neq 0$ and base it on the $x = x^{ss}$ and $\epsilon = 0$ situation, that is, the steady state, the true linear approximation of this value includes the $U_\epsilon^\alpha(x^{ss}, 0)\epsilon$ term. If the model has the

certainty-equivalence property then this term is zero; otherwise, this term is part of the linear approximation of $U^\alpha(x, \epsilon)$ based at $x = x^{**}$ and $\epsilon = 0$.

Differentiating (26) with respect to ϵ yields

$$0 = V_{i\epsilon} f^i + V_{ii} + \epsilon V_{iie} - \rho V_\epsilon \quad (31)$$

At the steady state of the deterministic case studied in the previous section, (31) reduces to

$$0 = V_{ii} - \rho V_\epsilon \quad (32)$$

which gives us our first correction, V_ϵ , term for variance.

We now move on to the other certainty nonequivalence correction terms. Differentiation of (31) with respect to x_j implies

$$0 = V_{ij\epsilon} f^i + V_{i\epsilon} f_j^i + V_{i\epsilon} f_\alpha^i U_j^\alpha + V_{iij} + \epsilon V_{iij\epsilon} - \rho V_{j\epsilon} \quad (33)$$

which at the steady state becomes

$$0 = V_{i\epsilon} f_j^i + V_{i\epsilon} f_\alpha^i U_j^\alpha + V_{iij} - \rho V_{j\epsilon} \quad (34)$$

which is a $n \times n$ linear system in the $V_{i\epsilon}$ unknowns. Differentiating (27) with respect to ϵ implies

$$0 = \pi_{\alpha\beta} U_\epsilon^\beta + V_{i\epsilon} f_\alpha^i + V_i f_{\alpha\beta}^i U_\epsilon^\beta \quad (35)$$

which, at the deterministic steady state, reduces to another linear system. Continued differentiation shows that computing higher-order terms also reduces to linear systems. Again, it is easy to compute the higher-order expansion terms of the Taylor series.

We have left out several details; in particular, we need to verify that these linear systems are nonsingular. Also, one needs proofs that the value functions are locally differentiable to the extent used in these formulas. These questions must be addressed on a problem-specific basis. If this approach leads to a singular system, then it breaks down, but that breakdown is checkable. Application of implicit function theorems should be able to prove if the value function is locally differentiable. What we have shown is that the standard linearization method frequently used in rational expectations analysis can be extended to include higher order terms and can be extended to model deviations from certainty equivalents, and that doing so will often be simple numerically.

2.5. Computational Costs of Linearization and Higher-Order Terms. The computations above have succeeded in showing us the qualitative nature of the problems which need to be solved. We next ask how large can we go with this approach. To see how large we can go, we need to know the timing demands of the basic operations in the procedure. These exercises were conducted using Matlab on a 90 Mz Pentium. The analysis of an n -state, m -control optimal control problem can be broken down into the following steps:

1. Compute steady state: time $\sim \mathcal{O}((2n + m)^3)$.
2. Compute Jacobian, A , of dynamic system at steady state: time $\sim \mathcal{O}((2n)^2)$.
3. Compute Jordan canonical form of $A = NDN^{-1}$: time $\sim \mathcal{O}((2n)^3)$.
4. Construct and solve the linear systems for higher-order and stochastic terms.

We now examine the computational cost of these steps. The computation of the steady state of an n -state, m -control problem is a nonlinear equation in $2n + m$ variables, which is not bad even for $n = m = 100$, and feasible for much larger problems. In fact, the steady state problem is often similar to CGE problems, hence we can use the special methods available for CGE models. Presumably this is not a difficult problem; if computing the steady state of the deterministic version is difficult, then it is unlikely that any method can produce a global solution to the stochastic version. Similarly, computing the Jacobian of the dynamic system is also presumably feasible. In general, the time demands of these steps varies greatly across problems, making it difficult for us to say anything generally. Hence, we will not consider these steps further, and instead focus on the generic features of this approach.

The critical limitations begin with the the eigenvalue-eigenvector extraction problem. This is a necessary step for any asymptotically valid linearization method. The limiting factor in computing the higher-order and stochastic terms is the cost of constructing and solving the linear systems which arise. The cost of constructing the linear systems depends on the specific problem and depends on the number of nonzero derivatives which arise in the systems (22), (23), and other equations, and the manner in which one computes those derivatives. We make no attempt to estimate the cost of

constructing these systems. We can give a reliable estimate of the cost of solving the resulting linear algebra problems. Table 1 contains the results of some experiments we ran to determine the cost of these steps. We took several random $n \times n$ matrices and computed eigenvalues and eigenvectors, for $n = 100, 200, 300, 400$. We next explored the timing for solving large linear systems of equations, the second operation. Table 1 also indicates the results for $n \times n$ matrices, $n = 100, 200, 500, 1000$. The third pair of columns in Table 1 display the mean time for several random matrix inversions; we do not report the variation in any column since there was little variation. The entries roughly reflect the order n^3 operation count associated with solving linear systems and inverting matrices.

To determine the cost of computing various order terms in the Taylor series expansion, we need to know the number of unknown coefficients of various orders. Table 2 indicates the size of the problem associated with various orders of the Taylor expansion for various dimensions; more precisely, it indicates the number of coefficients in the n 'th order terms of the Taylor expansion of a dimension d function. Note that the bottom right entry gives the general formula for the n 'th order coefficients in a d dimensional function.

These results give us good indications as to what is generally feasible. Since the linearization step involves computing a steady state, a Jacobian, and an eigenvalue-eigenvector decomposition, we see that moderately large systems can be linearized. The eigenvalue problem would generally be the most difficult, and we see here that a 10 state model would lead to a 20-dimensional eigenvector problem, one which takes under a second on a Pentium¹⁰. The computation of the quadratic terms is also possible. First we compute the V_{ij} terms using (19), a linear problem with, according to Table 2, 55 variables, and which can be solved using under .1 second¹¹. We then construct a linear problem involving the V_{ijm} , a problem with 220 variables, which takes roughly a second to solve. These values give us the information necessary to compute the U_{jm}^α , requiring 55 multiplications of 10×10 matrices. The third-order

terms of U need the 715 fourth-order terms of V from its defining linear system, a problem using about forty seconds, and then do 715 matrix multiplications, a problem taking less than three seconds according to Table 1. These results also show that quadratic approximations of control laws of 100-dimensional models is also feasible. Even cubic approximations could be had with patience and enough space. Furthermore, it is clear that the certainty nonequivalence terms we derived above could also be computed in the same time. Supercomputers could handle higher orders and/or larger models.

2.6. Global Accuracy. Perturbation methods produce the best possible asymptotically valid local approximations to a problem. However, we often want to use them for nonlocal approximations. We will not go into an extensive exploration of the global quality of the resulting approximations, a topic which must be investigated for specific problems and specific choices of tastes and technology. However, we should note that the existing literature is quite positive on the global quality of the resulting approximations. The Judd and Guu papers have investigated this issue in simple growth models. Typically, that is, for empirically reasonable choices of tastes and technology, they find that the linear approximations do well for small but nontrivial neighborhoods of the deterministic steady state and that the quality of the approximations improve substantially as the higher-order terms are added. They also find that the certainty nonequivalence terms are important to achieve high quality approximations for stochastic approximations. More precisely, they substitute the computed Taylor series into the defining equations and evaluate the resulting error. The resulting error for capital stocks near the steady state is often the order of machine zero, an accomplishment which few other methods can claim. While their investigations have been limited to relatively small models, there is no reason to suspect that the performance of this approach will decay drastically as we move to larger models. In any case, any user of these methods should use some diagnostics to estimate the region where the constructed series is a good approximation.

2.7. Summary of Perturbation Methods. The exercises above show that computing Taylor series expansions of dynamic optimization problems is feasible even for large problems, that is, problems with several states and controls. This includes

problems with several agents, several sectors, several goods, and several factors. Minor modifications allow these methods to be applied to models with distortions; see Judd(forthcoming) for discussions of how to do this. The method described above is a substantial improvement and generalization of the conventional linearization method used commonly in public finance and macroeconomics, allowing for certainty non-equivalence. In general, it can be used to create a linear theory for any observable quantity including those, such as risk premia, which depends on nonquadratic properties of tastes and technology.

The main disadvantage of this method is its local nature. We will next examine methods which are more global in their approach.

3. PROJECTION METHODS

Projection methods take a global approach to problems, making them applicable for a broader range of problems. However, they are much slower. Before discussing the large models we solve below, we shall review the basics of projection methods as applied to (1). That problem reduces to solving the functional equation

$$\begin{aligned} 0 &= u'(C(k, \theta)) - \beta E \{u'(C(F(k, \theta) - C(k, \theta), \tilde{\theta})) \\ &\quad \times F_k(F(k, \theta) - C(k, \theta), \tilde{\theta}) \mid \theta\} \\ &\equiv R(k, \theta; C) \end{aligned} \quad (36)$$

where $R(k, \theta; C)$ is the Euler residual of the policy function $C(k, \theta)$.

The projection approach to solving (36) can be broken down into four components. First, as noted above, we parameterize the unknown policy function and restrict it to lie in some finite-dimensional space, as in

$$\hat{C}(k, \theta; a) = \sum_{i=0}^n a_i \phi_i(k, \theta)$$

where the ϕ_i comprise a basis for all candidate functions. Second, we define a numerical approximation of the residual function applied to \hat{C} , $R(k, \theta; \hat{C})$; we let \hat{R} denote the numerical approximation of R . This primarily involves choosing a scheme for approximating the integral implicit in the conditional expectations. Third, we construct a collection of projection conditions; each is defined by some test function $\psi_\ell(k, \theta)$ relative to a weighting function $w(k, \theta)$, and ideally equals

$$\int R(k, \theta; \hat{C}(\cdot; a)) \psi_\ell(k, \theta) w(k, \theta) dk d\theta$$

but is implemented by choosing a finite sample (k_j, θ_j) , $j = 1, \dots, m$, and defining

$$P_\ell(a) = \sum_{j=1}^m \hat{R}(k_j, \theta_j; \hat{C}(\cdot; a)) \alpha_j \psi_\ell(k_j, \theta_j), \quad \ell = 1, \dots, n \quad (37)$$

Fourth, we choose some method to compute the coefficient vector a which solves the system $P_\ell(a) = 0$.

The various approaches to solving rational expectations models differ in their choices over these components. Table 3 displays the various choices. The critical fact is that this menu is “a la carte”: you make one choice from column A, then a choice from column B, etc., and almost any combination is possible. Table 3 outlines the basic components of projection methods, and points out the specific choices which have been and could be used.

The existing literature on computational rational expectations methods have pursued a small fraction of the possible combinations. Table 4 displays the various combinations which have been used. We left out the projection condition column since almost all essentially use collocation and Galerkin-style methods.

We will not consider the integration problem here, using product Gaussian quadrature in our examples; substantial improvement is possible by developing truly multidimensional quadrature and approximation methods, but we leave that for future research. In our examples we focus on some of the approximation choices and solution method choices. Judd(1992) contains the results for various methods for the representative agent model. We will now look at extensions of this model.

4. HETEROGENEOUS TASTES

We next examine how to introduce several agents into the analysis. We examine the same model as in (1) except we assume that there are n different types of agents, type i having utility function, $u_i(c)$, $i = 1, 2, \dots, n$, but a common discount factor β . In this case, the equilibrium decisions will depend on the distribution of wealth. Let $C^i(k)$ be

the consumption of type i agents when the wealth distribution is $k = (k_1, k_2, \dots, k_n)$. We assume that equity is the only asset which can be held. We do this to focus on the issues of size of models, whereas introducing an incomplete set of securities introduces other difficulties. The equilibrium of the resulting model is defined by the collection of Euler equations

$$R^i(k, \theta, C) = u'_i(C^i(k, \theta)) - \beta E \{u'_i(C^i(Y(k, \theta) - C(k, \theta), \tilde{\theta})) \times F_k(Y(k, \theta) - C(k, \theta), \tilde{\theta}) \mid \theta\}, \quad i = 1, 2, \dots, n$$

where $Y(k, \theta) \in R^n$ is the distribution of income in a period with initial capital stock distribution k and productivity θ , that is, $Y^i(k, \theta) = k_i F_1(k, \theta) + w(k, \theta)$ and $w(k, \theta) = F(k, \theta) - k F_1(k, \theta)$, where we use the tensor notation $k \equiv \sum_i k_i$. We focus on the residual functions

$$R^i(k, \theta, \hat{C}(\cdot; a)) = \hat{C}^i(k, \theta; a) - (u'_i)^{-1}(\beta E \{u'_i(\hat{C}^i(Y(k, \theta) - \hat{C}(k, \theta; a), \tilde{\theta}; a)) \times F_k((Y(k, \theta) - \hat{C}(k, \theta; a)), \tilde{\theta}) \mid \theta\}), \quad i = 1, 2, \dots, n$$

where we approximate each consumption function in terms of some unknown coefficients a , as in $\hat{C}^i(k, \theta; a)$. Due to the presence of the expectation operator, we need to form the approximate residual function for agent i ,

$$\hat{R}^i(k, \theta, \hat{C}(\cdot; a)) = \hat{C}^i(k, \theta; a) - (u'_i)^{-1}(\beta \hat{E} \{u'_i(\hat{C}^i(Y(k, \theta; a) - \hat{C}(k, \theta; a), \tilde{\theta}; a)) \times F_k((Y(k, \theta) - \hat{C}(k, \theta; a)), \tilde{\theta}) \mid \theta\}), \quad i = 1, 2, \dots, n$$

where \hat{E} represents some numerical approximation of the enclosed integral. In this paper we will use only product Gaussian quadrature formulas for integration. The identifying projections are

$$P_{ij}(a) \equiv \int_{\theta_m}^{\theta_M} \int_{k_m}^{k_M} \dots \int_{k_m}^{k_M} \hat{R}^i(k, \theta, \hat{C}(\cdot; a)) \psi_j(k, \theta) w(k, \theta) dk_1 \dots dk_n d\theta$$

where $i = 1, \dots, n$, and $j = 1, \dots, \hat{m}$. The computation of $P(a)$ also involves numerical integration; we will let $\hat{P}(a)$ denote a numerical integration approximation of $P(a)$; we will use product Gaussian quadrature here also; we will leave the integration issues for future research. The solution chooses a so that $\hat{P}(a) = 0$. We next discuss the leading possibilities for the approximation scheme and the solution method.

4.1. Representation: Tensor vs. Complete Polynomials. The tensor method approximates each consumption function as

$$\widehat{C}^i(k, \theta; a) = \sum_{j_1=0}^{n_k} \cdots \sum_{j_n=0}^{n_k} \sum_{\ell=0}^{n_\theta} a_{j_1 \dots j_n \ell}^i \varphi_{i_1}(k_1) \cdots \varphi_{i_n}(k_n) \psi_\ell(\theta), \quad i = 1, \dots, n$$

where $\varphi_i(k_j)$ ($\psi_\ell(\theta)$) is a degree $i-1$ ($\ell-1$) polynomial in k_j (θ) from some orthogonal family. We then solve for the unknown coefficients $a_{j_1 \dots j_n \ell}^i$. This is a method which can quickly get infeasible since the number of unknown coefficients equals $(n_\theta+1)(n_k+1)^n$ for each of the n policy functions. The complete polynomial method uses the form

$$C^i(k, \theta; a) = \sum_{\substack{0 \leq j_1 + \dots + j_n + \ell \leq d \\ 0 \leq j_i, \ell \leq d}} a_{j_1 \dots j_n \ell}^i \varphi_{j_1}(k_1) \cdots \varphi_{j_n}(k_n) \psi_\ell(\theta)$$

In this case, the total number of unknown coefficients is a far smaller number; in fact, the number of unknown coefficients here is the same as in the Taylor series expansion method, those numbers displayed in Table 3. In Table 3, we displayed the general formula for the number of coefficients. Note that it grows polynomially, not exponentially.

These general methods are two which are likely to be of general value. Other methods are not likely to be competitive for smooth models. For example, one would need far more unknown coefficients to use splines¹².

4.2. Solution: Successive Approximation vs. Newton's Method vs. Time Iteration. The next critical choice is the method we use to solve the projection equations for the coefficients a . Newton's method¹³ treats the conditions $P(a) = 0$ as a system of nonlinear equations and solves for a by repeated quadratic approximations. Newton's method is locally quadratically convergent, but each step uses $O(n^3)$ time because it computes a Jacobian. Some refinements economize on this by approximating the Jacobian, but the computational cost per step is still a problem.

Successive approximation proceeds more directly, uses less computation per step, but has only linear convergence if it converges at all. Specifically, successive approximation takes the policy functions computed in iteration j , $\widehat{C}^{i,j}$, and applies the computation

$$\widehat{C}^{i,j+1}(k, \theta) = (u')^{-1}(\beta \widehat{E} \{u'(\widehat{C}^{i,j}(Y(k, \theta) - \widehat{C}^j(k, \theta), \bar{\theta})) \times F_k(Y(k, \theta) - \widehat{C}^{i,j}(k, \theta), \bar{\theta}) \mid \theta)\}) \quad (38)$$

at a finite number of points (k, θ) to produce $\widehat{C}^{i,j+1}(k, \theta)$ data sufficient to fix the unknown coefficients of $\widehat{C}^{i,j+1}$. Since $\widehat{C}^{i,j+1}(k, \theta)$ is expressed directly in terms of the right hand side of (38), the computation cost is small. Successive approximation can be motivated by learning arguments in Marcet and Sargent(1989), but was actually used in the rational expectations literature earlier by Miranda and Helmburger(1988) who observed that it was an efficient method for computation.

Time iteration also uses the Euler equation to compute a new value for $\widehat{C}^{i,j+1}(k, \theta)$ but instead uses the equation

$$\begin{aligned} \widehat{C}^{i,j+1}(k, \theta) = & (u')^{-1}(\beta \widehat{E} \{u'_i(\widehat{C}^{i,j}(Y(k, \theta) - \widehat{C}^{i,j+1}(k, \theta), \tilde{\theta})) \\ & \times F_k(Y(k, \theta) - \widehat{C}^{i,j}(k, \theta), \tilde{\theta}) \mid \theta\}) \end{aligned} \quad (39)$$

is used to generate the necessary data. This is a much more complex way to fix $\widehat{C}^{i,j+1}(k, \theta)$ values since, for a fixed (k, θ) vector, (39) is a nonlinear equation in $\widehat{C}^{i,j+1}(k, \theta)$, hence involving more effort. Both successive approximation and time iteration are only linearly convergent. However, the computational demands of each iteration are only $O(n^2)$. Time iteration is more reliable but generally slower than successive approximation when the latter converges. Time iteration was used by Gustafson, and in the Wright and Williams work. In this paper, we will not discuss time iteration since experience¹⁴ and theory indicate that it will be much slower than Newton's method for small problems and slower than the successive approximation results below.

4.3. Accuracy. We need ways to ascertain if our solutions are "good." To measure the "accuracy" of our approximations we evaluate

$$E(a) \equiv \left\| R^i(\dots, \widehat{C}(\cdot; a)) / \widehat{C}^i(k, \theta; a) \right\|$$

at the solution for a for various norms, and use E as a unit-free measure of the "irrationality" in the approximate solution. For example, if the L_∞ norm is 10^{-2} then the Euler equation is accurate to within a penny per dollar of expenditure. Another way of expressing this is to say that the approximate policy functions $\widehat{C}^i(k, \theta; a)$ comprise an ϵ -equilibrium for $\epsilon = E$. This criterion is a strong one, much stronger than the one used in the Taylor-Uhlig symposium which essentially examined $E \{ R^i(\dots, \widehat{C}) \}$ where the expectation is taken over simulated paths for k and θ generated by \widehat{C} . The norm

we compute is nonzero as long as the Euler equation is nonzero anywhere, and the L_∞ norm is essentially the maximum Euler equation error over the region explored. We parameterize the production function so that the symmetric deterministic steady state capital stock is 1 for each agent, and then let $k_m = .5$ and $k_M = 1.5$, a large range.

4.4. Results. We now give results for these alternative approaches. In Table 5 we examine 2, 3, 4, and 5 agent models using both tensor product and complete polynomial bases, and using both Newton's method and successive approximation. In all cases we use product Gaussian quadrature to compute all integrals and we use a simple linear initial guess which goes through the deterministic steady state and zero. The column under γ lists the different values of relative risk aversion for the different agents; these values cover the range generally considered reasonable. We converged when coefficients were deemed within 10^{-5} of the solution. The accuracy is the base 10 logarithm of the L_2 definition of E above applied to the worst agents Euler equation; in fact, the Euler equation errors were all very close for all agents. The magnitude of the accuracy measure is therefore (roughly speaking) the decimal digit accuracy. These examples were computed on a 50MHz 486 machine.

The results are along expected lines. First, the total amount of time for Newton's method is roughly cubic in the number of unknown coefficients. Newton's method is generally more accurate than successive approximation, but for large problems takes much more time. The other problem is that storing the Jacobian can demand more space than available in RAM; this happens for the larger models. Note that the complete polynomial method is generally efficient but not tensor product methods.

The successive approximation method turns out to dominate Newton's method for large models. In fact, for the higher-order approximations of the five-agent model, only successive approximation with complete polynomials is feasible given space requirements. The successive approximation method turns out to be stable (given the initial condition), a fortunate property of these models. Judd (forthcoming) shows that successive approximation applied to the single-agent, deterministic model can be

unstable when the elasticity of substitution is large, but is stable near the deterministic steady state for most reasonable choices of γ . This stability property apparently is robust to multiple agent models.

5. HETEROGENEOUS WEALTH, COMMON TASTES

A simpler case is where all agents have the same intertemporal tastes, but may have different wealth. Here we can exploit obvious symmetry properties to drastically reduce the computational burden. We now assume that the n different agents have the same utility function, $u(c)$, and a common discount factor β . Outside of some special cases, we cannot aggregate preferences to reduce the model to a representative agent. Let $C^i(k)$ be the consumption of type i agents when the wealth distribution is $k = (k_1, k_2, \dots, k_n)$, where k_i is the capital owned by a type i agent. The equilibrium is again defined by the collection of Euler equations

$$u'(C^i(k, \theta)) = \beta E \{u'_i(C^i(Y(k, \theta) - C(k, \theta), \tilde{\theta})) F_1(Y(k, \theta) - C^i(k, \theta), \tilde{\theta}) \mid \theta\} \quad (40)$$

for $i = 1, 2, \dots, n$, where $Y(k, \theta) \in R^n$ is the distribution of income in a period with initial capital stock k and productivity θ , that is, $Y^i(k, \theta) = k_i F_1(k, \theta) + w(k, \theta)$. The tensor method approximates each consumption function as

$$C^\ell(k, \theta; \mathbf{a}) = \sum_{i_1=0}^{n_k} \cdots \sum_{i_m=0}^{n_k} \sum_{j=0}^{n_\theta} a_{i_1 \dots i_m j}^\ell \varphi_{i_1}(k_1) \cdots \varphi_{i_m}(k_m) \psi_j(\theta)$$

where $\varphi_i(k_m)$ ($\psi_j(\theta)$) is a degree $i-1$ ($j-1$) polynomial in k_m (θ) from some orthogonal family. We then solve for the unknown coefficients $a_{i_1 \dots i_m j}^\ell$. However, we can impose certain symmetry conditions. First, an agent's behavior depends only on his wealth and the distribution of wealth, and all agents have the same consumption function. Therefore, if $i_\ell = i_{\ell'}$ then the coefficients $a_{i_1 \dots i_\ell \dots i_m j}^\ell = a_{i_1 \dots i_{\ell'} \dots i_m j}^{\ell'}$ for all ℓ and ℓ' . Furthermore, $a_{i_1 \dots i_\ell \dots i_m j}^1 = a_{i_1 \pi(i_2) \dots \pi(i_\ell) \dots \pi(i_m) j}^1$ for all permutations π on $\{i_2, i_3, \dots, i_m\}$. Together, these conditions imply that the policy functions are of the form

$$C(k, \theta; \mathbf{a}) = \sum_{i,j} a_{ij} \varphi_i(k_1) \Psi_j(k_2, k_3, \dots, k_n)$$

where $\Psi_j(k_2, k_3, \dots, k_n)$ is a symmetric polynomial in the $n-1$ variables k_2, k_3, \dots, k_n . The degree 1, 2, and 3 symmetric polynomials are listed below in Table 6:

The complete polynomial method uses, for a type 1 agent, the form

$$C^1(k, \theta; \mathbf{a}) = \sum_{0 \leq \deg(\phi_i) + \deg(\Psi_j) + \deg(\psi_\ell) \leq n} a_{ij\ell} \varphi_i(k_1) \Psi_j(k_2, k_3, \dots, k_n) \psi_\ell(\theta)$$

where the consumption functions of the other types are constructed by symmetry considerations. To identify the unknown coefficients, we use the identifying projections

$$0 = \int \dots \int u'(C^1(k, \theta)) - \beta E \{u'(C^1(Y(k, \theta) - C(k, \theta), \tilde{\theta})) \\ \times F_k(Y(k, \theta) - C(k, \theta), \tilde{\theta}) \mid \theta\} \varphi_i(k_1) \Psi_j(k_2, k_3, \dots, k_n) \psi_\ell(\theta) dk d\theta$$

for i, j , and ℓ such that $0 \leq \deg(\phi_i) + \deg(\Psi_j) + \deg(\psi_\ell) \leq n$. In this case, the total number of unknown coefficients is a far smaller number than the unrestricted case, and there is only one Euler equation which needs to be fitted. The result is a far smaller system than we would have if we had followed the general multi-agent approach of the previous section. To save on space, we do not discuss computational results since they are exactly what one would expect: imposing the symmetry conditions results in far faster computations without any loss of accuracy.

6. CONCLUSION

This paper has shown that it is feasible to compute rational expectations models substantially more complex than the usual representative agent, single good model. Using only 486- and Pentium-class personal computers, we have shown how to use perturbation methods to solve dynamic models with up to 50 state variables (that is, 50 agents or 50 capital stocks, or some combination) in only minutes. We have also demonstrated that more globally oriented methods can produce accurate approximations of multiple-agent models.

We also have seen that the methods of choice depend on the size of the problem. Table 7 summarizes our findings. Our experiments indicate that the dominant approach to large models with Euler equation formulations will be successive approximation solution methods combined with complete polynomial bases, outperforming Newton-based methods and monotone- and contraction-operator methods. The apparent dominance of successive approximation indicates that we need to better un-

derstand the stability problems of successive approximation and develop approaches to detect and deal with them.

This work clearly indicates that rational expectations models of moderate size can be reliably and quickly solved numerically. This paper focussed on the approximation and solution methods which have been used, and used only the simplest integration methods. Exploitation of more advanced approximation, solution, and integration techniques will surely lead to drastic improvements. We also suspect that these methods are capable of making efficient use of supercomputing environments, making even larger models feasible. Overall, given the limited use of available hardware and software in this paper, we believe that numerical solution of large stochastic, dynamic models is an attainable goal.

REFERENCES

- [1] Bensoussan, Alain. *Perturbation Methods in Optimal Control*, John Wiley and Sons, 1988.
- [2] Bizer, David S., and Kenneth L. Judd. 1989. "Uncertainty and Taxation", *American Economic Review*, 79:331-336.
- [3] Blanchard, Olivier Jean and Charles M. Kahn. 1980. "The Solution of Linear Difference Models under Rational Expectations" *Econometrica* 48:1305-1311.
- [4] Brock, William A., and L. J. Mirman. 1972. "Optimal Economic Growth and Uncertainty: The Discounted Case," *Journal of Economic Theory* 4: 479-513.
- [5] Brock, William A., and Stephen J. Turnovsky. 1981. "The Analysis of Macroeconomic Policies in Perfect Foresight Equilibrium," *International Economic Review* 22:179-209.
- [6] Christiano, Lawrence J. 1990. "Solving the Stochastic Growth Model by Linear-Quadratic Approximation and by Value-Function Iteration," *Journal of Business and Economic Statistics*, 8:23-26.
- [7] W. J. Coleman, II. 1990. "Solving the Stochastic Growth Model by Policy Function Iteration," *Journal of Business and Economic Statistics*, 8:27-29.
- [8] Cooley, Thomas F., and Gary D. Hansen. 1989. "The Inflation Tax in a Real Business Cycle Model." *American Economic Review*.
- [9] Daniel, J. W. 1976. "Splines and Efficiency in Dynamic Programming," *Journal of Mathematical Analysis and Applications* 54:402-407.
- [10] Dantzig, George B., R. P. Harvey, Z. F. Landowne, and R. D. McKnight. *DYGAM- A Computer System for the Solution of Dynamic Programs*, Control Analysis Corporation, Palo Alto, California. 1974.
- [11] Fischer, Stanley. "Capital Accumulation on the Transition Path in a Monetary Optimizing Model." 1979. *Econometrica* 47:1433-39.

- [12] Fleming, Wendell. 1971. "Stochastic Control for Small Noise Intensities," *SIAM Journal of Control* 9 (No. 3, August, 1971).
- [13] Robert L. Gustafson, "Carryover Levels for Grains: A Method for Determining Amounts That are Optimal Under Specified Conditions," USDA Technical Bulletin 1178, 1958.
- [14] den Haan, Wouter, and Albert Marcet. 1990. "Solving the Stochastic Growth Model by Parameterizing Expectations," *Journal of Business and Economic Statistics*, 8:31-34
- [15] Hall, Robert. 1971. "The Dynamic Effects of Fiscal Policy in an Economy with Foresight," *Review of Economic Studies* 38:229-244.
- [16] Judd, Kenneth L. 1982. "An Alternative to Steady-State Comparisons in Perfect Foresight Models," *Economics Letters* 10:55-59.
- [17] Judd, Kenneth L. 1985. "Short-run Analysis of Fiscal Policy in a Simple Perfect Foresight Model", *Journal of Political Economy* 93:298-319.
- [18] Judd, Kenneth L. 1992. "Projection Methods for Solving Aggregate Growth Models", *Journal of Economic Theory* 58:410-452.
- [19] Judd, Kenneth L. *Numerical Methods in Economics*, MIT Press (forthcoming, 1997).
- [20] Judd, Kenneth L. 1996. "Approximation, Perturbation, and Projection Methods in Economic Analysis," in *Handbook of Computational Economics*, edited by Hans Amman, David Kendrick, and John Rust, North Holland.
- [21] Judd, Kenneth L., and Jess Gaspar. "Perturbation Methods for Discrete-Time Dynamic Deterministic Models," Hoover Institution, mimeo, 1997.
- [22] Judd, Kenneth L., and Sy-Ming Guu. 1993. "Perturbation Solution Methods for Economic Growth Models," in *Economic and Financial Modeling with Mathematica*, Hal Varian, ed., Springer-Verlag: New York.

- [23] Judd, Kenneth L., and Sy-Ming Guu. "Asymptotic Methods for Aggregate Growth Models," forthcoming in *Journal of Economic Dynamics and Control*.
- [24] Kydland, Finn E., and Edward C. Prescott. 1982. "Time to Build and Aggregate Fluctuations," *Econometrica* 50: 1345-1370.
- [25] Laitner, John. 1981. "The Stability of Steady States in Perfect Foresight Models," *Econometrica* 49:319-333.
- [26] Laitner, John. 1984. "Transition Time Paths for Overlapping-Generations Models," *Journal of Economic Dynamics and Control* 7:111-129.
- [27] Magill, J. P. Michael. 1977. "A Local Analysis of N-Sector Capital Accumulation under Uncertainty," *Journal of Economic Theory*, 15: 211-219.
- [28] Marcet, Albert. "Simulation Analysis of Dynamic Stochastic Models: Applications to Theory and Estimation," in C. Sims (ed.) *Advances in Econometrics Sixth World Congress*, Vol II., 91-118, Cambridge University Press.
- [29] Marcet, Albert, and Thomas J. Sargent. 1989. "Convergence of Least Squares Learning Mechanisms in Self-referential Linear Stochastic Models," *Journal of Economic Theory* 48:337-68.
- [30] McGrattan, Ellen R. 1990. "Solving The Stochastic Growth Model by Linear-Quadratic Approximation." *Journal of Business and Economic Statistics* 8: 41-43.
- [31] Miranda, Mario J., and Peter G. Helmberger. 1988. "The Effects of Commodity Price Stabilization Programs." *American Economic Review* 78:46-58.
- [32] Prescott, Edward C., and R. Mehra. 1978. "Recursive Competitive Equilibrium: The Case of Homogeneous Agents," *Econometrica* 48:1365-1379.
- [33] Stokey, Nancy L., and Robert E. Lucas, Jr. *Recursive Methods in Economic Dynamics*, Harvard University Press: Cambridge. 1989.

- [34] Taylor, John B., and Harald Uhlig. 1990. "Solving Nonlinear Stochastic Growth Models: A Comparison of Alternative Solution Methods," *Journal of Business and Economic Statistics*, 8:1-18.
- [35] Wright, Brian D., and Jeffrey C. Williams. 1984. "The Welfare Effects of the Introduction of Storage," *Quarterly Journal of Economics* 99:169-182.
- [36] Williams, Jeffrey, and Brian Wright, *Storage and Commodity Markets*, Cambridge University Press, January, 1991.

FOOTNOTES

*The authors acknowledge the comments of Wouter den Haan, two anonymous referees, and conference participants. All remaining errors are the responsibility of the authors. Dr. Judd also acknowledges the financial support of NSF grant SBR-9309613.

¹The term "projection method" is a catchall term in the mathematical literature which includes "method of weighted residuals," "finite element," "boundary element," "Galerkin," "least squares," "Rayleigh-Ritz," and other similar methods.

²Wright and Williams(1984) and some later writers parameterize a conditional expectation function, such as the one on the right hand side of (2), which characterizes the solution. If one is to use polynomial approximation methods, then one should approximate some continuous function which characterizes equilibrium. Wright and Williams show that when price and policy functions have kinks, it is better to approximate the conditional expectations function. Our comments apply equally to this case since, as is clear from Judd(1992), the key fact is that one is solving for an unknown smooth function.

³For example, the procedure we outline below is *not* the procedure discussed by McGrattan; in general, our Taylor series expansion method will produce different results.

⁴By "correct," I refer to the concepts, definitions, methods, and solutions derived in the mathematical literature on the stable manifold theorem and asymptotics. The interested reader should consult Bensoussan, Fleming, Fleming and Souganides, and Coddington and Levinson. Judd(forthcoming) gives an example of where one proposed method goes wrong, and a more extensive discussion of these points.

⁵Marcet did not indicate what he meant by "perturbation" methods. It is true that some of the "linear approximation" methods discussed in the macroeconomic literature are not easily extended to compute higher-order terms. However, it has always been known in the mathematical literature that higher-order terms can easily be computed if one used what the mathematical literature calls perturbation methods; see Bensoussan and the extensive citations there, for example.

⁶This is also the method used in dynamic analyses of the 1970's and early 1980's, such as Hall(1971), Fischer(1979), Judd(1982), Laitner(1984), and many others. Many

later writers reject the standard approach in favor of ad hoc procedures, but never explain why. For example, Cooley and Hansen's(1989) analysis of monetary equilibria near the deterministic steady state ignores the simpler procedure outlined in Fischer.

⁷Taylor series expansions are valid over nontrivial intervals for analytic functions, but we make no claim that our problems have analytic solutions.

⁸By a "linear theory of u " we mean an approximation of u which is linear in the state variables x , and which is asymptotically valid as x converges to the steady state value of x .

⁹The discrete-time approach can be similarly analyzed, but at a greater notational cost.

¹⁰This estimate is based on log interpolation of the $n = 10$ and $n = 100$ cases in the "Eigenvector" results in Table 1.

¹¹Again, this is based on log interpolating in the "Linear System" column in Table 1 between the $n = 10$ and $n = 100$ cases.

¹²One possible alternative is to create a problem specific basis, an approach defined and discussed in Judd(1996) and papers cited there.

¹³We actually use an implementation of Powell's hybrid method. Our experience is that Powell's method is far faster than the pure Newton method. We have tried a few other methods on various projection problems but only with disappointing results.

¹⁴For example, compare the execution times in Coleman's implementation of time iteration and the Newton method results reported in Judd(1992). The experience with time iteration in Bizer and Judd(1989) also was disappointing.

TABLES

Table 1: Linear Algebra Times

Eigenvectors:		Linear Systems:		Matrix Inversion:	
<i>n</i>	seconds:	<i>n</i>	seconds:	<i>n</i>	seconds:
10	.1	10	.0025	10	.05
100	2.6	100	.17	100	.27
200	25.5	200	.77	200	2.4
300	104	500	18.3	300	9.0
400	230	1000	150	400	24.4

Table 2: Taylor Series

dimension	5					10			
order	1	2	3	4	5	1	2	3	4
# coefficients	5	15	35	70	126	10	55	220	715
dimension	20					50			
order	1	2	3	4		1	2		
# coefficients	20	210	1540	8855		50	1275		
dimension	100					<i>d</i>			
order	1	2	3			<i>n</i>			
# coefficients	100	5050	171700				$\frac{(n+d)!}{n!d!}$	$-\frac{(n+d-1)!}{(n-1)!d!}$	

Table 3: Projection Method Components

Approximation	Integration	Projection Conditions	Solution Method
Piecewise Linear	Newton-Cotes	Galerkin	Newton
Ordinary Poly.	Gaussian Quad.	Collocation	S.A.-time iteration
Orthogonal Poly.	Monte Carlo	Sum of Squares	S.A.-learning
Splines	quasi-Monte Carlo	Subdomain	Least. Squares
Neural Networks	R^n Rules	Method of Moments	Global Minimum
Finite Element	Derivative Rules		Homotopy
Customized Bases	Asymptotics		

Table 4: Choices Made in Literature

Authors	Approximation	Integration	Sol'n Method
Gustafson(1959)	piecewise linear	Newt.-Cotes	S.A.-time it.
Wright-W.(1982,4)	polynomial (of cond. exp.)	Newt.-Cotes	S.A.-time it.
Miranda-H.(1986)	polynomials	Newt.-Cotes	S.A.-learning
Bizer-Judd(1989)	piecewise linear	Newt.-Cotes	S.A.-time it.
Coleman(1990)	finite element	Gaussian	S.A.-time it.
den Haan-M.(1990)	polynomial (of cond. exp.)	Sim. M.C.	S.A.-learning
Judd(1992)	orthogonal polynomial	Gaussian	Newton

Table 5: Time and Accuracy Comparisons

number agents	tastes (γ)	degree	basis	number coef's	Newton's time	Method: accuracy	Successive time	Approx: accuracy
1	-2	1	t	4	:0.05	-2.7	:0.22	-2.7
			c	3	:0.06	-2.6	:0.39	-2.6
		2	t	9	:0.22	-3.4	:0.55	-3.4
			c	6	:0.17	-3.3	:0.93	-3.3
		3	t	16	:0.71	-4.1	:1.15	-4.1
			c	10	:0.49	-4.0	:1.92	-4.0
		4	t	25	:1.7	-4.8	:2.15	-4.9
			c	30	:0.99	-4.7	:3.29	-4.6
2	-1.1	1	t	16	:0.66	-3.1	:1.49	-3.1
			c	6	:0.38	-2.7	:1.42	-2.7
	2	t	54	:7.3	-4.1	:8.02	-4.1	
		c	20	:2.5	-3.4	:6.43	-3.4	
	3	t	128	1:22	-5.0	:32.9	-4.5	
		c	40	11.4	-4.1	:20.8	-4.1	
	4	t	250	12:34	-5.9	1:48	-4.5	
		c	70	:45.2	-4.8	:55.5	-4.7	

Table 5: Time and Accuracy Comparisons (Continued)

number agents	tastes (γ)	degree	basis	number coef's	Newton's time	Method: accuracy	Successive time	Approx: accuracy
3	-1.1	1	t	48	:6.93	-3.4	:7.5	-3.4
			c	15	1.48	-2.8	:4.6	-2.8
	-2	2	t	243	7:07	-4.6	2:11	-4.5
			c	63	:20.8	-3.6	:36.4	-3.6
	-3	3	t	768	inf	inf	19:57	-4.6
			c	105	4:05	-4.3	3:09	-4.3
	4	t	1875	inf	inf	1 hr 56	-4.6	
			c	210	46:58	-4.9	12:45	-4.8
4	-1.1	1	t	128	1:09	-3.5	:33.1	-3.5
			c	24	:5.10	-2.9	:13.3	-2.9
	-2	2	t	972	inf	inf	24:57	-4.6
			c	84	2:47	-3.7	3:04	-3.7
	-3	3	t	4096	inf	inf	7 hr 13	-4.6
			c	224	52:11	-4.4	26:01	-4.4
5	-1.1	1	t	320	8:52	-3.6	2:48	-3.6
			c	35	:17.90	-3.0	:38	-3.0
	-2	2	t	3645	inf	inf	5 hr 16	-4.6
			c	140	12:18	-3.8	10:18	-3.8
	-3	3	t	20,480	inf	inf	inf	inf
			c	420	12 hr 50	-4.5	3 hr 27	-4.5

Note: "inf" means infeasible. "h hrs n : m.l" means "h hours n minutes, m.l seconds".

Table 6: Symmetric Polynomials

Degree:	Polynomials:		
1	$x + y + \dots + z$		
2	$x^2 + y^2 + \dots + z^2$	$(x + y + \dots + z)^2$	
3	$x^3 + y^3 + \dots + z^3$	$x^2y + x^2z + \dots + y^2z + \dots$	$(x + y + \dots + z)^3$

Table 7: Final Comparisons

Method:	Basis:	Solution Method:	Advantages:	Disadvantages:
Taylor Series	Complete	Eigenvalues, linear equations	Very fast	Local validity
Projection methods	Tensor or complete	Newton	Quadratic convergence	Infeasible for large problems
	Tensor or complete	Successive approximation	Easy Iterations	possible nonconvergence